
2014 FISHING FORECAST AND TIPS



Welcome to the 2014 Fishing Forecast for Kentucky's major fisheries. The forecast is based on 2013 fish population surveys, creel surveys, fish stockings, and historical knowledge of the fisheries. This handout is designed to assist anglers in planning their fishing trips and improving their fishing success. Additional fishing information is available from the Department's website at fw.ky.gov or by obtaining copies of the 2014 Sport Fishing and Boating Guide available at most sporting goods stores. The Kentucky Trout Waters brochure is contained in the 2014 Fishing and Boating Guide.

To locate fishing access sites in Kentucky, visit our website and click on "Fish" and then "Where to Fish". You will be able to search for your favorite water bodies and get directions to all major boat launches and access sites.

Opening day of the 2014 fishing season starts March 1 with the new year's license, so take a trip to your local sporting good store, get online at fw.ky.gov, or call 1-877-598-2401 to purchase your 2014 fishing license.

The Fishing Forecast was partially financed through funds provided by your purchase of fishing equipment and motor boat fuels under the Federal Sport Fish Restoration Program.





New and Expanding Fisheries in 2014

- Rainbow trout will be stocked in Morton's Lake (Higginson-Henry WMA; Union County), Flycatcher Lake (Peabody WMA; Ohio, Hopkins and Muhlenberg counties) and Sandy Lee Watkins Park Lake (Henderson County). See the Trout Waters section of the 2014 Fishing and Boating Guide for specific stocking months.
- Brown trout will be stocked in Jennings Creek (Warren County), Roundstone Creek (Hart County),
 Sulphur Springs Creek (Simpson County) and the Nolin Tailwater (Edmonson County).
- With completion of the work on Wolf Creek Dam, rainbow trout stockings in the Cumberland Tailwater will be increased in anticipation of better water quality.
- Saugeye have been stocked into Guist Creek, Bullock Pen, A.J. Jolly and Boltz lakes to create an additional fishery.
- Sauger stockings continue in the Salt, Green and Barren rivers with expanded fishing opportunities expected below Taylorsville, Green and Barren dams.

Fishing Forecast Cheat Sheet

Make sure you check out the new Fishing Forecast Cheat Sheet located at the end of this publication. For those looking for a quick answer to where the best fishing can be found for each species, the Cheat Sheet is your guide. The Cheat Sheet lets you know which lakes are forecasted to provide good to excellent fishing for each of your favorite fish species. Just look up the species you are interested in and then see which lakes have a check mark next to them. These are the lakes which will provide the best chance for a memorable fishing trip for that species. The Cheat Sheet is located on two pages, so if you don't see the species you are interested in on the first page; make sure to check the following page.

New Up-and-Comers

Several lakes and rivers have shown improvements in their fisheries in the last year. You might try and get in on the action at the following water bodies:

- Largemouth bass on the Ohio River (multiple counties).
- Bluegill at Lake Reba (Madison County) and Mill Creek Lake (Monroe County).
- Redear sunfish at Cedar Creek Lake (Lincoln County)
- Rainbow trout on the Cumberland Tailwater (Russell, Cumberland, Clinton and Monroe counties).
- Brown trout on the Cumberland Tailwater (Russell, Cumberland, Clinton and Monroe counties).
- Striped bass at Lake Cumberland (Russell, Wayne, Clinton and Pulaski counties).
- White bass at Barkley Lake (Trigg, Lyon and Livingston counties).
- Channel catfish at Lake Malone (Muhlenberg, Todd and Logan counties).
- Blue catfish at Dewey Lake (Floyd County).

Late Winter/Early Spring Fishing Frenzy

As warm winds begin to flow over Kentucky, put your new fishing license to good use by trying a few early spring fishing hot spots. In March, fish activity begins in the smaller bodies of water such as farm ponds and smaller public lakes as water temperatures reach the 50's. These smaller, shallower bodies of water warm quicker than reservoirs in spring. Late winter and early spring may be the best time of the year to catch big fish.

Trophy largemouth bass, especially females, are at their heaviest weight of the entire year during late winter and early spring. Female bass feed heavily during this time to provide fuel for the upcoming spawn. Farm ponds and public lakes are prime spots for early spring largemouth bass fishing before the larger bodies of water warm enough for bass to feed.

A few of the best lakes for catching trophy bass in spring are Lake Barkley, Kentucky Lake and Lake Malone in western Kentucky along with Cedar Creek, Kincaid and Guist Creek lakes in central Kentucky. Greenbo Lake in northeast Kentucky and Wood Creek Lake in southeast Kentucky both produced multiple state record largemouth bass in the past. Woods Creek Lake is the home of the current 13-pound, 10.4 ounce state record. Barren River Lake in west-central Kentucky, Lake Beshear and Lake Mauzy in west Kentucky along with Lake Herrington and Green River lakes in central Kentucky also offer excellent early spring bass fishing. Dewey Lake, Fishtrap and Fishpond lakes give east Kentucky anglers a great shot for an early spring trophy largemouth bass.

If you are after smallmouth bass, head to Green River, Kentucky, Cumberland, Laurel River, Dale Hollow and Fishtrap lakes for a good chance at catching the smallmouth bass of a lifetime in late winter and early spring. Streams also provide excellent fishing for smallmouth bass in early spring throughout central and eastern Kentucky. Adult fish migrate from their winter habitat in deep, slow moving holes in the lower sections of streams to spawning areas in smaller tributary streams. Elkhorn Creek, South Fork, Licking River and several upper Kentucky, Green and Barren River tributaries offer the best spring fishing opportunities for smallmouth bass.

Slab crappie are ready to be caught, beginning in late winter, at many reservoirs in central and western Kentucky. Catches of crappie over 12 inches are not uncommon at Barkley and Kentucky lakes during this time. Black crappie move into shallow water well before the spawning season in early March, so start early for these fish. Other good bets for crappie include Herrington, Cumberland, Dewey, Rough, Nolin, Barren and Buckhorn lakes as well as Ohio River embayments.

Excellent white bass fishing can be found during the early spring in the upper sections of reservoirs such as Barkley, Cave Run, Nolin River and Herrington lakes. The Kentucky, Green and Ohio rivers offer excellent spring white bass fishing.

During late winter, sauger and walleye fishing peaks in tailwaters below dams on the Ohio, lower Kentucky, Cumberland and Tennessee rivers. Lake Cumberland, Laurel River and Carr Creek lakes possess excellent walleye fisheries.

If you are interested in muskellunge, they move in spring to shallow shoreline structure and headwater areas in search of warmer water at Cave Run, Green, and Buckhorn lakes. Remember, a 36-inch minimum size limit and one fish daily creel limit is in effect for muskellunge at these three reservoirs.

Do not forget to take advantage of trout waters such as seasonal catch-and-release trout streams, lakes stocked with trout in winter months, Fishing in Neighborhood (FINs) lakes stocked with trout and the Cumberland River tailwater below Lake Cumberland in late winter and early spring. Trout bite willingly in cold water. Expect fishing to improve on the Cumberland Tailwater now that the construction on Wolf Creek Dam is completed.

Try these late winter and early spring hot spots. Not only will they cure your cabin fever, but you might catch the fish of your lifetime.

2014 FISHING FORECAST

Kentucky Department of Fish and Wildlife Resources

The fishing forecast is based on 2013 fish population surveys, planned fish stockings, and projections for 2014.

Location/Species	Forecast	Comments
BARKLEY LAKE and	TAILWATER	45,600 acres (Trigg, Lyon and Livingston counties)
Blue catfish	Excellent	Good numbers of 10-20 pounders with the occasional 50+ pounder. Best fishing is during late spring and summer along deep river channels. Look for good flow in the lake for best fishing conditions. Blue catfish can be caught year-round in the tailwater.
Bluegill	Excellent	Good numbers of fish up to 9 inches; the best fishing comes in May on sand or gravel spawning beds. Crickets and worms under corks during the spawn is a great option. Fish move deeper as the water warms in summer.
Channel catfish	Excellent	Best fishing is during May along rocky banks and during summer along the main river channel and adjacent flats. Try stink-bait in shallow water in the spring. Channel catfish can also be caught year-round in the tailwater.
Crappie	Good	Good numbers of keeper-size fish in the 10-12 inch size range. Many fishermen are changing tactics, casting or trolling light jigs and roadrunners year round.
Largemouth bass	Excellent	Very good fishing with excellent numbers of >15-inch fish, and numbers of >20-inch fish are well above average. Shallow fishing for spawning bass can be amazing in the spring around button ball bushes with floating baits and plastic worms. Post spawn fish move to drop offs in the main lake and mouths of the bays. Try fishing Carolina rigs, jigs and crankbaits in deeper water. In fall and winter, bass move into the bays to feed on shad.
Redear sunfish	Excellent	Good numbers of 8-10 inch fish with trophies up to 12 inches. Prime fishing occurs in late April and throughout May as fish are spawning. Fish can be found in areas with gravel substrate and vegetation. Redear generally spawn a bit deeper than bluegill.
Sauger	Poor	Population consisting of low numbers of harvestable-size (>14-inch) fish. Many fish are caught trolling deep diving crankbaits near secondary channel breaks during the early summer. Sauger can also be caught while fishing for catfish along the main river channel. Fair fishing in the tailwaters.
Smallmouth bass	Fair	Smallmouth bass make up a small portion of the total bass population but are common in certain habitats. Fishing is best around rocky shorelines near deep water, main lake points, and the canal during late winter and early spring with suspending jerk baits.
Striped bass	Good	The tailwater fishing is best for consistent catches, but the lake produces good catches as well. Fish deep main channel ledges near bridges and other places with current for best luck trolling crankbaits or fishing vertical with large jigs. Good trophy potential.
White bass	Good	Fishing success has been decent the last few years. Look for schools of shad breaking the surface during summer and fall. If you can't find them working the surface, fish deeper along ledges on the main lake with a spoon or inline spinner. Similar to striped bass, good fishing occurs in the tailwaters.

_ocation/Species	Forecast	Comments
BARREN RIVER (from	Barren River La	ke dam to Green River confluence)
Hybrid striped bass	Fair/Good	Barren River Lake tailwater area hybrid fishing is best, especially during or following periods of heavy lake discharges. Good fishing also below instream dams at Bowling Green and Greencastle.
Muskellunge	Good	Target mouth of feeder creeks and around wood structure or other current breaks from dam to Green River confluence. Stocked annually.
Smallmouth bass	Good/Excellent	Good numbers of 12-inch fish. Trophy fish (20-inch plus) available. Fishing best from dam to mouth of Drakes Creek. Target areas above and below riffle/shoal areas and rocky banks in pool areas.
Spotted bass	Fair	Found throughout Barren River. Fish slower moving pools with rocky banks and logs. Fish spinner baits and crankbaits in the spring and early summer. Buzzbaits during th summer.
Walleye/Sauger	Fair/Good	Fishing best below instream dams (Barren River Lake, Greencastle and Bowling Green) especially during December – March. Good size walleye (15-25 inch) availabl Sauger have been stocked since 2010.

ARREN RIVER LAKE	10,000 acres	(Allen and Barren counties)
Summertime (mid Ju	une through Septe	ember) oxygen levels are too low to support fish below 15-18 feet.
Bluegill (sunfish)	Good	Good numbers of 6–8 inch fish. Fish stump beds and back of coves during the spring and fall. Fish deeper in the summer near cove mouths and along bluff walls near laydowns or standing timber.
Catfish	Excellent	Good populations of channel and flathead catfish present with good numbers of larger individuals. Fishing best near creek channels and at night. Blue catfish have been stocked annually since 2010, with some of these already over 20 inches. Fish for blues with live or cut bait in open water areas with jugs or rod and reel.
Crappie	Good/Excellent	Good numbers of larger crappie (10-11 inches; black and white) available; however, for best numbers of whites, fish winter and early spring. Crappie mix has shifted to about 60-40 ratio (black crappie to white crappie). Good numbers of both are available due to good year classes/spawns of both in 2008 and 2010 and also 2011 for black crappie. Target shallower water to find black crappie in spring and early summer. Summer and winter fishing is best around deeper structure, fish attractor/brush sites or near creek channel.
Hybrid striped bass	Good	Overall numbers are good with large fish (20-inch plus) available. Spring fishing best in headwater areas with spinners, crank baits and curly tailed jigs. Target main lake (around islands in mid-lower lake) early in the morning and at dusk in the summer and fall as schools appear throughout the lake. Use crankbaits and spoons. 200,000 stocked annually since 2000. Should see extra bump in numbers of 11-16 inch fish from elevated stocking (400,000 fish) in 2013.
Largemouth bass	Excellent	Good numbers overall and good numbers of 15-inch plus and 20-inch plus fish.
Smallmouth bass	Fair	Low density population overall, but quality/trophy fish (20-inch plus) available. Target lower half of the lake near the dam, fishing rock outcroppings and rocky transition areas.
Spotted bass	Fair	Lower density population overall, but quality fish (15-inch plus) available. Bridge pylons and bluff walls with rock slides are good places to target spots.

_ocation/Species	Forecast	Comments
BARREN RIVER LA	KE (continued	I)
White bass	Fair	Overall numbers remain low. Best opportunity is during spawning run in late March to mid April with in-line spinners, small jigs and small crankbaits in Barren River, Peter, Skaggs and Beaver Creeks above the lake.
Yellow bass	Excellent	Abundant and willing to bite. Most fish are 7-9 inches in length. Bite readily on small jigs, spinners and minnows. Bridge pylons excellent areas to target for summer time bite.
BEAVER LAKE	158 acres	(Anderson County)
Bluegill	Good	High numbers of 6-8 inch fish; ≥ 8.0 inch fish present.
Channel catfish	Good	Around 3,700 catchable-size fish stocked each year; fish over 15 inches common. Fa fishing good in shallows in coves.
Crappie	Fair/Good	Small population; good numbers of fish between 8-10 inches; larger fish possible.
Largemouth bass	Good	Good numbers of bass up to 15 inches; improved numbers of fish ≥15 inches.
Redear sunfish	Good	Many 7-9 inch fish available around vegetation; good numbers of 10-11 inch fish possible.
BESHEAR LAKE	700	(Coldwall and Christian acception)
		(Caldwell and Christian counties)
Blue catfish	Fair	Blue catfish have been stocked every other year since 2005. Their population is doing fair, with good growth. Some already measure over 25 inches and weigh over 5 pounds.
Bluegill	Fair	Numerous small fish with occasional 8-inch fish caught.
Channel catfish	Excellent	High numbers of 10-15 inch channel catfish in the population. Channel catfish are stocked every other year at 10-12 inches. This is an underutilized fishery. Most anglers fishing for catfish use jug lines or noodles. Catfish are one of the reasons people go to Lake Beshear to fish.
Crappie	Fair	Numerous fish in the 6-8 inch range, with an occasional 10 incher. Best fishing during spring in areas of aquatic vegetation and around woody structure. The population is considered stunted; therefore it is ok to harvest the small crappie.
Largemouth bass	Excellent	One of the main reason anglers fish Lake Beshear. The bass population is dominate by 12-18 inch largemouth bass. Best fishing occurs during spring when fish move to shallow water along the water willow and lily pads. Try fishing the edges of these

Location/Species	Forecast	Comments
BOLTZ LAKE 92 a	acres (Grant (County)
Blue catfish	Fair	Catchable-size fish stocked each year; 25-inch plus fish possible.
Bluegill	Good	Good numbers of fish between 6-8 inches.
Channel catfish	Fair	Around 1,700 catchable-size fish stocked each year.
Crappie	Fair	Abundant; majority of fish between 6-8 inches; larger fish possible due to recent removal of smaller crappie in spring of 2012.
Largemouth bass	Fair	Good numbers bass at and above size limit (12-inch size limit); some potential for bass ≥20 inches. A little over 2,100 6-inch bass were stocked in spring of 2013.
Redear sunfish	Poor	Low numbers of 6-8 inch fish available around vegetation.
BRIGGS LAKE 18	acres (Logai	n County)
Summertime (mid-		ptember) oxygen levels are too low to support fish below 7 feet.
Bluegill	Good	Good numbers of 6-inch fish available.
Channel catfish	Good	All sizes available. Stocked annually.
Crappie	Fair	Large fish present; fish laydown trees, stakebeds and brushpiles in late winter and early spring.
Largemouth bass	Fair	Better than average numbers of 12-inch plus fish.
Redear sunfish	Fair	Good numbers of larger fish (8-inch plus) available. Larger fish very shallow in early spring; then move to thermocline depth.
BUCKHORN LAKE a	nd TAILWATER	1,250 acres (Leslie and Perry counties)
Bluegill	Good	Good size distribution and very good numbers. Upper lake areas and heads of coves hold best numbers during summer pool. Shallow mudflats in main lake areas can hold large congregations of fish on spawning beds.
Catfish	Excellent	High numbers of channel catfish with good recruitment of young fish each year. Some flathead catfish available up to 40-50 pounds. Tailwater provides additional good fishery.
Crappie	Fair	Large number of fish just below the 9-inch minimum size limit. If the majority of these fish surpass the 9-inch size in 2014 there will be a lot of keeper fish available. July and August can provide some very good fishing over shallow mud flats adjacent to main lake channel. Try trolling bait tipped jigs or small crankbaits over the flats. Also, late September/early October during drawdown to winter pool can be very good in the same locations.
Largemouth bass	Fair	Excellent growth rates of young bass and fair numbers of 15-inch plus fish; also good numbers for fish greater than 20 inches. Spring fishing success can be very good if high water events do not occur.

ocation/Species	Forecast	Comments
BUCKHORN LAKE an	d TAILWATER (d	continued)
Muskellunge	Good	Fish numbers over 36 inches are starting to increase and fish numbers overall are good. Fishing can be difficult for periods following flood events. January-March fish will congregate in the lower lake and near dam. Trace Branch area has good fishing in July and August. September will find many large fish in backs of coves in shallow water along edges of weedbeds. Bank fishing opportunities are good at the tailwater in March.
White bass	Fair	Some successful recruitment of young fish in 2013 will increase numbers for 2014. Most fish will be made up of 2 year classes measuring 9-10 and 13-15 inches.
BULLOCK PEN LAKE	134 acres	(Grant County)
Blue catfish	Fair/Good	Catchable-size fish stocked each year; 25-inch plus fish possible.
Bluegill	Poor	Most fish are 6 inches or less, but are abundant.
Channel catfish	Fair	Catchable-size fish stocked each year; potential for larger fish.
Crappie	Fair	Abundant population of small fish.
Largemouth bass	Good/Excellent	Good numbers of bass at and above size limit (12-inch size limit); excellent potential for quality-size fish and bass over 20 inches.
Redear sunfish	Poor	Very low numbers.
White bass	Poor/Fair	Small population of large fish in the 14-17 inch range.
CARPENTER LAKE	68 acres (Da	viess County)
	Fair/Good	Good number of 6 to 7-inch bluegill, but few over 8 inches.
Bluegill	raii/Guuu	Good number of 6 to 7-inch bluegill, but lew over 6 inches.
Channel catfish	Good	Stocked annually. Excellent growth rates produce good numbers of 15-20 inch and some larger fish.
Largemouth bass	Fair/Good	Increasing number of sublegal (8-11 inch) fish in the population, but the number of 15-20 inch fish is decent and holding steady.
Redear sunfish	Fair	Most redear 7-8 inches with a few in the 9-11 inch range.

Hydrilla was documented in Carr Creek Lake for the first time in 2008. Boaters are asked to take measures to prevent the spread of this damaging non-native invasive species within the lake and between lakes by properly inspecting and disinfecting boats, trailers and equipment.

Bluegill	Fair	Good numbers of small bluegill present. Shallow water areas in creek arms provide good fishing in spring and early summer. Fishing pier near ramp at marina holds fish in submerged brush.
Crappie	Fair	Larger black crappie from 9-12 inches and white crappie from 10-16 inches. Deadfall trees and submerged brush piles will congregate large numbers of fish during most of the year. Research project with white crappie is ongoing.
Largemouth bass	Fair	Good numbers of large fish over 20 inches. March-early April anglers have very good fishing success for big fish. Usually poor recruitment of small fish and stocking of fingerlings occurs each year.
Smallmouth bass	Poor	Low numbers but fish sampled up to 20 inches.
Spotted bass	Excellent	Best numbers are concentrated in lower and mid lake; a few large fish up to 18 inches; excellent numbers through 16 inches.
Walleye	Excellent	Very high numbers of fish from 15-26 inches; a few fish up to 28 inches; March through May are good months to try during day or night. Standing timber will hold suspended fish in summer as well as open water areas with deeper schools of alewife that walleye will suspend beneath.

CAVE RUN LAKE 8,270 acres (Bath, Menifee, Rowan and Morgan counties)

The beds of aquatic vegetation composed of Eurasian milfoil and curly pondweed can still be found in many areas of the lake. Barring extremely high water conditions in 2014, these weed beds should continue to provide good largemouth bass and muskie fishing locations.

Bluegill	Good	Widespread within the lake; most 7-inches long or less, but it is not unusual to catch 7 to 8 inch long bluegill. Look for bluegill early in the year in the back of coves or other shallow water areas that warm up first or where they may spawn. Later in the year, bluegill can be found around stumps or sunken woody structure within many areas of the lake. Fishing main lake structure during the summer can yield good results.
Catfish	Good	Good numbers of channel catfish up to two feet long with a few larger fish. Many are found along shallow rocky shoreline areas in early May-June where they seek cover in rock piles and submerged logs to spawn. Later they tend to be found in deeper water. Although not as plentiful, some trophy-sized flathead catfish are found in the lake.
Crappie	Good	During 2013 anglers continued to have very good success catching crappie. The lake still holds nice numbers of 2 to 3 year old fish and they should continue to push through the population maintaining the good numbers of larger fish for a few more years. During spring months look for crappie in the shallows near shore and later in the year you can find black crappie in association with aquatic vegetation and white crappie around deeper structure.

Location/Species	Forecast	Comments
CAVE RUN LAKE (co	ntinued)	
Largemouth bass	Excellent	The North Fork and Licking River portions of the lake (upper reaches) generally hold the larger fish. In these areas try fishing the rocky banks warmed by the morning sun in April and May. For the rest of the lake fish the edges and pockets of the weed beds, fallen trees and woody structure within embayments. Electrofishing catch rates for fish over 15 inches and over 20 inches were the highest ever recorded on Cave Run Lake in 2013. This should continue into 2014 and should provide excellent fishing for larger fish. Note: 13-16 inch protective slot limit.
Muskellunge	Excellent	Excellent angling can still be expected in this fishery. Trophy fish can be found primarily in the spawning month of April in the backs of the large coves or in the fall in association with gentle sloping gravely banks or near channel drop offs. General fall fishing for muskellunge can also be hot from the first part of September until the end of October in the shallow flats area and in the upper reaches of the lake. Note: 36 inch minimum size limit, one fish daily creel limit.
Smallmouth bass	Fair	While still uncommon, smallmouth have been picking up in recent years. They are found primarily in the lower lake (mouths of Licking River and Beaver Creek to the Dam). Look for them off rocky points or shoreline areas along the main lake. One excellent bank on the lake is directly across from the Twin Knobs beach area. Other areas to try include: Scotts Creek, rocky areas around the confluence of the Licking River, and the rocky bank from Scotts Creek Boat Ramp to the dam. Note: 18-inch minimum size limit.
Spotted bass	Good	Numerous within the middle and lower main lake shoreline; become less common within the upper reaches of the lake. Most less than 13-inches long.
White bass	Good	A white bass die off in 2013 leaves some question marks about the strength of the population. Certainly, not all fish vanished with this kill, but it did put a hit on the larger fish, especially in the main lake. These fish will begin their rebound in the next few years. Note: no more than 5 white bass in the daily creel limit may be greater than 15-inches long.
CEDAR CREEK LAKE	784 acres	(Lincoln County)
Bluegill	Good	Excellent numbers of bluegill in lake, although most are less than 7 inches. Bank anglers can choose from several designated fishing access areas for family fishing.
Channel catfish	Fair	Fair numbers of 12-18 inch fish with a few over 20 inches. We have reduced the stocking rate the last few years due to slower growth of stocked fish.
Crappie	Good	The crappie are growing better in recent years and there are currently good numbers of mostly moderate-sized (9-10 inch) crappie with a few large ones.
Largemouth bass	Excellent	Wide range of sizes available for anglers, including an impressive number of fish larger than 18 inches. Cedar Creek Lake is a great place to catch a large bass. Note: a trophy regulation of a 20-inch minimum size limit, 1 fish daily creel limit, is in effect on this lake.
Rainbow trout	Fair	Stocked in late fall and early spring to provide a cool weather fishery.
Redear sunfish	Good	Good numbers of 7-10 inch fish available. The number of larger redear sunfish has improved in recent years. Try mealworms around vegetation for best fishing.

ocation/Species	Forecast	Comments
CLEAR CREEK LAKE	40 acres	(Bath County)
	in 2007- 2009; p	a good portion of the lake; therefore fishing may be difficult especially in the fall. Grass clease release any that are caught. Due to federal ownership of this lake, the use of his time.
Bluegill	Good	Good numbers of 6 to 7-inch bluegill are present with occasional 8 inch fish caught. Fish shallow shoreline areas in the spring especially behind standing timber.
Largemouth bass	Fair	Lake receives heavy pressure reducing the overall numbers of bass, but a few in the 20-25 inch range have been sampled.
Redear sunfish	Good	Good population of 7-8 inch fish. Fish shallow shoreline areas in the spring especia behind standing timber.
CORINTH LAKE 96	6 acres (Grar	nt County)
Bluegill	Fair	Abundant; majority of fish 6 inches or less. Shad removal conducted in 2013 should improve bluegill population.
Channel catfish	Good	Catchable-size fish stocked each year; fish over 15 inches common.
Crappie	Fair	Few fish; most between 8 to 9 inches.
Largemouth bass	Good	Good numbers of bass at and above size limit (12-inch size limit); potential for bass between 15-20 inches.
Redear sunfish	Good	Good numbers of fish greater than 7 inches; Numbers of quality-size fish should increase due to shad removal.
CUMBERLAND LAKE	50,250 acre	es (Russell, Wayne, Clinton, and Pulaski counties)
The boundaries of L	_ake Cumberlan	nd are now defined as Cumberland Falls on the Cumberland River, Devils Jump on the Eastle River and Laurel River Lake Dam on Laurel River.
Bluegill	Fair	Fair numbers of 6-7 inch bluegill in lake, with some larger fish available. Try crickets coves and cuts for good early summer action.
Channel catfish	Excellent	Outlook remains bright with excellent numbers of 16-20 inch fish. Nightcrawlers still fished or slowly trolled on main lake flats and cuts during late summer or fall should
		pick up some quality fish.
Crappie	Good	· · · · · · · · · · · · · · · · · · ·
Crappie Largemouth bass	Good	pick up some quality fish. Population remains good with some quality crappie (12-inch plus) available. Newly flooded willows and sycamore trees should benefit the crappie population for the foreseeable future. Try jigs or minnows in headwaters of the lake (near mouth of Laurel and Rockcastle Rivers) in early spring for some excellent fishing. Fall fishing should also be good in major tributaries in the upper half of lake. Try jigs or minnow
		pick up some quality fish. Population remains good with some quality crappie (12-inch plus) available. Newly flooded willows and sycamore trees should benefit the crappie population for the foreseeable future. Try jigs or minnows in headwaters of the lake (near mouth of Laurel and Rockcastle Rivers) in early spring for some excellent fishing. Fall fishing should also be good in major tributaries in the upper half of lake. Try jigs or minnow around brush near heads of creeks for best results. Stable density of 14-16 inch fish; some larger fish are available. Creek arms in upper

CUMBERLAND LAKE	E (continued)		
Striped bass	Good	The striped bass population is improving and anglers can expect to catch slightly larger fish in 2014. The 2009 year class (mostly 24-28 inch fish) should provide the bulk of the striped bass fishery but some larger fish are in the population. The lake received additional fish in 2013, in anticipation of full lake levels in 2014. The 22-inch size limit remains in effect, and anglers should have no trouble catching their limit.	
Walleye	Fair/Good	Anglers can expect a modest boost in walleye size and numbers this year, as the walleye population continues to rebound from lows in 2011 and 2012. Expect most fish to average 17-22 inches, with a few over 4 pounds. Stocking numbers were increased in 2013 in anticipation of full lake levels in 2014. Best fishing in spring is at night on surface baits. Night crawler rigs and deep diving crank baits trolled 20-35 feet deep along the main lake shoreline work well during summer and fall.	
White bass	Fair	Numbers remain low but some good-size whites are available. Upper half of lake is best for numbers. For best success, try headwater areas in early spring.	
CUMBERLAND LAKE	F TAII WATER	(Russell, Cumberland, Clinton, and Monroe counties)	
Brook trout	Fair	Annual stockings have established brook trout in the tailwater. Most fish range from 10 13 inches, but some larger fish are present. The potential for breaking the state record exists in this relatively new fishery.	
Brown trout	Good	Numbers and size of brown trout remain significantly decreased but are improving. Anglers should find good numbers of small browns and fair numbers of keeper fish (>20 inches). Try fishing at night with large crankbaits or spinners to target bigger fish.	
Rainbow trout	Good	Rainbow trout abundance is still significantly lower than peak years, but nearing the long term average. Adequate numbers of stocked fish should be available to anglers and larger fish (>15 inches) were boosted by additional stockings in 2013. Expect fishing to continue to improve as flows return to normal.	
Striped bass	Fair	The tailwater provides the best location in state to catch a trophy striper. Fairly low density population but good size (20 lbs +). Mid and lower sections of river are best. Try using live bait or trolling shad colored crankbaits or swim baits.	
Walleye/Sauger	Fair	Decent population of walleye in tailwater, but tends to fluctuate yearly. Try near dam in late winter and early spring with small jigs and grubs. From mid-summer to fall, fish below Burkesville trolling shad colored crankbaits or nightcrawler rigs. Moderate population of sauger with quality fish (2-5 lb).	
White bass	Fair	Population appears to have leveled off. Try casting spinners and jigs near the dam or near creek mouths in the lower section in early spring for best success.	
DALE HOLLOW LAK	E 4,300 acre	es in Kentucky (Clinton and Cumberland counties)	
Largemouth bass	Good	Good size distribution with moderate numbers over 15 inches.	
Smallmouth bass	Good	Continues to produce impressive numbers of fish over 3 lbs. Standard fare of jigs and soft plastics work well as do float-n-fly techniques in cold water months.	
Spotted bass	Good	Good density of quality-size fish (12-14 inches).	

Location/Specie	s Forecast	Comments
DEWEY LAKE	1,100 acres (Floyd	County)

Zebra mussels and hydrilla are present at Dewey Lake. Boaters are asked to take measures to prevent the spread of these damaging non-native invasive species by properly inspecting and disinfecting boats, trailers and equipment.

Note: New regulations for blue and channel catfish beginning March 1, 2014. 15-fish daily creel limit; only 1 fish in the daily creel may be longer than 25 inches.

the daily creet may	y be longer man z	to money.
Blue catfish	Good	Blues have been stocked annually from 2009 through 2012 to start this fishery. Population is represented well by all year classes. Larger fish up to 30 inches.
Bluegill	Good	Stable population with good growth. Large fish are around 7 inches long.
Channel catfish	Excellent	Good numbers found in the lower lake during winter months. The upper lake is best in the summer months. Lots of good bank fishing access.
Crappie	Excellent	Crappie have shown good growth in recent years. Another year with good catches of 9- 12 inch crappie is expected in 2014. Black and white crappie are present with larger white crappie to 15 inches and black crappie to 11 inches.
Flathead catfish	Good	Good size distribution; a few large fish from 20-40 pounds caught each year; popular with ticklers/noodlers in June; try fishing shallow in September and October.
Largemouth bass	Fair	Bass continue to show good size distribution. The 2009 and 2010 year classes are strong. There were high numbers of 13 and 14 inch fish in 2012. Many of these fish will be legal-size in 2014.
Redear sunfish	Fair	Fish are present through 12 inches. Some areas such as the rip-rap rock face of dam hold good fish. Increasing population of zebra mussels will improve this fishery. Try fishing bait near the bottom around weedy areas or brush. If catching bluegill, then go deeper or further off shoreline to get into redear sunfish.
White bass	Fair	Numbers remain stable; good spawns in 2008, 2010 and 2012. John's Creek will congregate fish in March for spawning (look for Redbud trees in bloom).
X RIVER – below H	errington Lake (Ga	arrard and Mercer counties)
Brown Trout	Fair	Currently no holdover fish due to continuously high water conditions. One stocking (usually June) per year.
Rainbow Trout	Fair	Currently no holdover fish due to continuously high water conditions. Stocked frequently throughout the year. Use artificial lures or flies from Herrington Lake Dam downstream for 2 miles.
KHORN CREEK –	main stem and No	orth Fork (Franklin and Scott counties)
Channel catfish	Good	Large fish possible in bigger pools. Good numbers of "keeper" fish, especially in the lower section of the main stem.
Largemouth bass	Fair/Good	Quality fishing found in slow moving pools in the North Fork. Note: 12-16 inch slot limit applies on main stem.
Rock Bass	Fair	Many fish between 7-9 inches in main stem and in the lower North Fork section.
Smallmouth bass	Good/Excellent	Very good numbers of smallmouth bass, especially on the main stem within the slot. Also good numbers of fish greater than 16 inches. Note: fish from 12-16 inches (slot

limit) are protected in the main stem Elkhorn Creek.

Location/Species	Forecast	Comments
ELMER DAVIS LAKE	149 acres (Owen County)
Bluegill	Fair	Decent numbers of fish between 6-8 inches; good numbers found around beaver lodges. Bluegill population is being effected by presence of gizzard shad in the lake.
Channel catfish	Good	Around 3,200 stocked each year; fish over 15 inches possible.
Crappie	Fair	Most around 9-11 inches.
Largemouth bass	Good/Excellent	Plenty of bass under the 12-15 inch slot limit with good numbers in the protected slot. Bass numbers greater than 15 inches have increased slightly.
Redear sunfish	Good	Fish greater than 8 inches possible near weed beds or on spawning beds in upper embayments. Population is being negatively effected by shad.
Warmouth	Fair	Nice-size fish present, usually around weed beds. Locally known as "redeye" or "rockbass".

FAGAN BRANCH LAKE 140 acres (Marion County)

Summertime (mid-June through September) temperature and oxygen levels for trout are best in the 20-30 foot range. This zone decreases as the summer progresses. Lake is extremely clear year round.

Bluegill	Fair	Fair numbers of 7-inch plus fish available.
Channel catfish	Good	Good numbers of 11-14 inch fish. Stocked every other year.
Crappie	Fair	Best fishing early and late and at night under floating lights using minnows. Fair numbers of fish, but not many larger fish (10-inch plus).
Largemouth bass	Fair	Stable population but slower growing. Fair numbers of 12-inch plus fish available.
Rainbow trout	Fair	Best fishing will be at night under floating lights from May–October due to clear water. Fish will be in 20-30 foot ranges during the summer. Fish available throughout the lake during the winter. Stocked semi-annually (November and February) since 2010.
Redear sunfish	Good/Excellent	Above average numbers of larger fish (8-inch plus) present. Fish edges of deep weed beds (15-20 feet) during summer.

Location/Species	Forecast	Comments
FISHPOND LAKE	32 acres	(Letcher County)
Bluegill	Good	Population is doing well with fish common up to 10 inches; largest individuals will be deeper than young fish; try night fishing for large fish under overhanging shoreline brush and trees; warmouth are another panfish that are available at Fishpond and reach large sizes there.
Channel catfish	Fair	Lake receives annual stocking; fish up to 21 inches present; night fishing may produce best results.
Largemouth bass	Excellent	Very good size distribution of fish up to 24 inches and 10 pounds; water is very clear and fish are often very difficult to catch during daylight hours. When rainbow trout are stocked, they are heavily targeted by bass.
Rainbow trout	Good	Fishing is best around the months when fish are stocked; stockings are in January, April, May and October. Due to clear water, most fishing is done at night. However, when a new stocking has just occurred, the daytime fishing can also be good. A few trout will oversummer in most years providing fish up to 16 inches.

FISHTRAP LAKE and TAILWATER 1,131 acres (Pike County)

Zebra mussels are present at Fishtrap Lake. Boaters are asked to take measures to prevent the spread of this damaging non-native invasive species by properly inspecting and disinfecting boats, trailers and equipment.

Note: new regulations for blue and channel catfish beginning March 1, 2014. 15-fish daily creel limit; only 1 fish in the daily creel may be longer than 25 inches.

Blue catfish	Fair	Blues have been stocked in 2011 and 2013 to start this fishery for use of zebra mussels. Larger fish are 20 inches. Fishery should continue to improve with increasing population of zebra mussels and large gizzard shad population.
Bluegill	Excellent	Some large bluegill available up to 11-12 inches; May and June have been popular months for angling.
Channel catfish	Excellent	Good size distribution and numbers found throughout the lake and into Levisa Fork.
Crappie	Good	High numbers of 8 inch fish in fall 2013 samples. This will provide a lot of fish over the 9 inch length limit in spring of 2014. Size distribution and numbers are good through 12 inches with some larger fish available to 14-15 inches. This fishery consists only of white crappie.
Flathead catfish	Excellent	Numerous large fish with good numbers found in the lake and into Levisa Fork. Numerous rocky ledges and banks around the lake provide excellent noodling opportunities when fish are spawning.
Hybrid striped bass	Excellent	Very good numbers of hybrids in the lower lake in late summer and at winter pool; fair spawning run in headwaters in April; night fishing has become popular; look for fish in the jumps from May-July; good distribution of fish through 10 pounds. This is an underutilized fishery.
Largemouth bass	Good	Fair numbers with excellent size distribution and good growth; lake is prone to rapid changes in water level affecting fish activity for periods, ratio of keeper size bass is one of the best in eastern Kentucky. Fingerlings were stocked in 2010, 2012 and 2013 to supplement those year classes in the lake.
Redear sunfish	Fair	This fishery was initiated with stockings from 2010-2013. There were some 9 and 10 inch fish in angler catches during 2012 and 2013.
Smallmouth bass	Good	Make up approximately 15% of black bass; good fishing primarily in the headwaters of the lake up into the Levisa Fork River and in the lower portion of the lake from Grapevine to dam; large individuals up to 22 inches. Approximately 30,000 stocked in 2010. Excellent numbers in the Levisa Fork upstream of the lake.

Location/Species	Forecast	Comments
Floyds Fork Creek	27 miles (Jeff	erson County)
Largemouth bass	Poor	Small percentage of all three black bass species. Most under 10 inches. Note: new regulation ; 15-inch minimum size limit on largemouth and smallmouth bass; 1 fish daily creel limit; beginning March 2014.
Rainbow trout	Good	Fish stocked 3 times per year, mainly 8-12 inches.
Rock bass	Fair	Most fish between 6-8 inches. More numerous in the middle section of the stream.
Smallmouth bass	Good	Good numbers of smallmouth bass 9-12 inches. Note: new regulation; 15-inch minimum size limit on largemouth and smallmouth bass; 1 fish daily creel limit; beginning March 2014.
GRAYSON LAKE	1,512 acres (C	Carter and Elliott counties)
Bluegill	Good	Numerous, up to 7 inches long. Widespread throughout the lake as well as other kinds of panfish. Try fishing 5-6 feet deep in May and June.
Catfish	Good	Widespread throughout the lake. Fair numbers of channel and flathead catfish up to 20-inches long; some trophy-size flathead catfish available. Try fishing the middle and upper sections in early fall.
Crappie	Fair	Population samples collected in 2012 were far better than previous years. For this fish it is important to find cover or fish deep near channel edges around standing timber.
Hybrid striped bass	: Fair	Hybrid striped bass were originally stocked from 1998-2003 and this program resumed in 2009. Fish from the original stocking can still be found and range in length from 24 - 27 inches. Good numbers of fish from the most recent stocking program in the ≥15 inch range have been sampled.
Largemouth bass	Fair	Fair numbers of fish over 15 inches, but most still less than the minimum size limit. Most prevalent of all black bass species in the lake. Better population exists in the upper reaches than the lower reaches. High water helps to congregate these fish within the weeds and shoreline vegetation. Note: 15 inch minimum size limit.
Spotted bass	Fair	Stable population. Most fish are found in lower 2/3 of lake.
Smallmouth bass	Poor	Rarely found while sampling. The best bet is the lower portion of the lake. Note: 15-inch minimum size limit.
GREEN RIVER (da	am at Pochastar	down to mouth at Spottsville)
Catfish	Good	Good populations of channel, blue and flathead catfish. Trophy size blue and flathead
Callisti	Good	catfish available. Good bets are tailwaters below locks and dams, in deeper holes, and around logs, brush, or rocks.
Crappie	Fair/Good	Fish tributary mouths, below locks and dams, and brush along bank in spring and fall with live minnows.
Striped and hybrid striped bass	Good	Stripers and hybrids congregate below dams late spring through early fall. Cast top water plugs or lead head jigs with plastic bodies into boils below dams. Shad imitation swim baits and bucktail jigs also produce well.
White bass	Good/Excellen	t Fish mouth of tributary streams in spring and below dams mid-spring through fall. Use top-water baits or lead head jigs below dams and lead head jigs or live bait at mouth of tributary streams during spring. Fish bank access area and boat ramp at mouth of Pond River late March through April.

Location/Species	Forecast	Comments
GREEN RIVER (dar	m at Rochester up	to Green River Lake Dam)
Muskellunge	Good	Fishing best at feeder creek mouths and around woody debris or other current breaks. Fishing below instream dams also good bet. Stocked annually throughout river above Rochester.
Rock bass	Excellent	Good numbers of 6-8 inch fish present from dam to Munfordville.
Smallmouth bass	Good	Good numbers overall with larger fish (15-inch plus) available. Fishing best from Greek River Lake dam to Mammoth Cave.
Walleye	Fair	Fishing best below Green River Lake dam area and instream dams during late winter and early spring.
GREEN RIVER LAKE	8,210 acres	(Taylor and Adair counties)
	,	October) oxygen levels are too low to support fish below 18-23 feet.
`	J	, , , , , , , , , , , , , , , , , , , ,
Bluegill	Fair/Good	Large population with occasional 7-8 inch fish. Fish stump beds and back ends of coves during spring and deep-water brush, standing timber and structure during summer for larger fish.
Channel catfish	Good/Excellent	All sizes available. Hook and line fishing best at night or when water levels are on the rise and muddy. Jug fishing good mid-summer to fall.
Largemouth bass	Excellent	Exceptional numbers of quality-size fish (15-inches plus) and better than average numbers of 20-inch plus fish.
Muskellunge	Excellent	Legal-size fish (36-inch) abundant. Good numbers of 40-inch plus fish. Fall, late winter and spring excellent time to catch trophy fish. Population maintained by annual stocking.
Smallmouth bass	Fair	Low density overall, but can expect a slight upturn in numbers from good spawn of 2010. Expect better than average numbers of 15-inch plus fish. Fish in mid-lower sections of Robinson Creek and Green River arms for best numbers and chance for 20 inch plus fish.
Spotted bass	Good	Better than average numbers of 12-inch plus fish. 15-inch plus spots not uncommon.
Walleye	Fair	Overall numbers are fair as are numbers of larger fish (20-inch plus). Fish dam area and headwater areas of Robinson Creek and Green River arms in late February through mid to late March. Target standing timber coves, main lake stump beds, humps, and channel drop offs during summer and fall. Population maintained by annual stocking.
White bass	Poor	Overall numbers are low. Department is in process of stocking the lake to re-establish population.
White crappie	Good/Excellent	Good numbers of harvestable-size fish (9-inch plus). Expect to weed through lots of sub-legal fish to find keeper fish.

Location/Species	Forecast	Comments
GREENBO LAKE	181 acres (0	Greenup County - Greenbo State Park)
Bluegill	Fair	Gradually improving population; larger fish (up to 10 inches long) are caught early to late spring in deep-water structure. Try fishing near beaver lodges. Note: bluegill and other sunfish - 15 fish daily creel limit.
Channel catfish	Good	Stocked annually; many 2-3 pound fish observed during fall sampling. Note: limb lines (set lines), jug lines and trot lines are illegal to use at this lake.
Largemouth bass	Good	Good bass population but very difficult lake to fish; good numbers of fish over 15-inches long; potential for trophy-size bass continues. Due to water clarity, anglers may have to fish deeper when fishing this lake. Based on 2013 spring sampling data, anglers should notice their catch of fish in the 12-14 inch size range increase.
Rainbow trout	Excellent	Annually stocked population with some carry-over; best fishing year round is in the open lake and near the dam; during the spring and summer try fishing at night under lights. Winter fishing can also be good.
Redear sunfish	Fair	Numbers low, however a few 8 to 13-inch fish continue to be caught. After the spawn try fishing in deeper water near the bottom. Note: see bluegill regulation.
GUIST CREEK LAKE	317 acres	(Shelby County)
Bluegill	Fair	Abundant, most less than 6 inches. Easy to catch; good place to take a child fishing
Diacym	i dii	with live bait.
Channel catfish	Excellent	Catchable-size fish stocked each year; very good numbers of fish with potential for trophies. Note: this lake is home to current bullhead catfish and white catfish state records.
Common carp	Good	Bottom-fish with grain-based bait; 3 to 4-lb. fish common.
Crappie	Fair	Most in the 7-10 inch range.
Hybrid striped bass	s Fair	Small population; good numbers of fish in the 16 to 22-inch range. Around 19,000 1.5-inch fish were stocked in June 2013.
Largemouth bass	Excellent	Excellent numbers of fish over the 12-inch size limit; excellent potential for a trophy fish
HERRINGTON LAKE	<u> </u>	•
Bluegill	Good	Stable number of fish up to 8 inches in and around cover in embayments and inlets of rock walls; best April-June.
Common carp	Fair	Fish in coves or headwaters with grain-based baits on bottom.
Crappie	Good	Difficult to locate, but many quality-size fish (9 inches or larger) available. Best around brush in upper half of lake. Potential for large black crappie around debris in inlets in main lake.
Flathead catfish	Good	All sizes present; usually feed at night on sunfish and shad.
Hybrid striped bass	s Fair	Two to five pound fish are common; possible trophy-size fish.
Largemouth bass	Good	Good numbers of fish over 12 inches, many over 15 inches. Largest fish around shoreline cover in March and April; potential for trophy-size fish. Best success in the upper half of lake. Good night fishing during the summer.

Location/Species	Forecast	Comments	
HERRINGTON LAKE (continued)			
Longnose gar	Good	Large fish common; look for them around schools of shad. Schooling common in fall; potential trophy size.	
White bass	Fair	Should see good numbers during the spring run up the Dix River. Fair numbers of fish in the 12-14 inch range Winter and spring rains should improve the spring spawning run.	
KENTUCKY LAKE ar	nd TAILWATER	51,000 acres (Marshall, Calloway, Lyon, Livingston and Trigg counties)	
Blue catfish	Excellent	In the lake, best fishing is along the main river channel where the deeper drops are at. Fish congregate along the ledges of the channel when there is current and scatter along the adjacent flats during periods of low flow. Drift using live shiners or cut shad/skipjack in areas where creek channels meet the main river channel and along the main river channel near bends. Use a depth finder to determine if the fish are near the bottom or suspended and then fish accordingly. Other baits to try include chicken hearts, night crawlers, grass hoppers or catalpa worms.	
		In the tailwaters there is year-round fishing. The ideal times are in April and May, and October and November. Best to use fresh bait (shad and skipjack herring) caught in the tailwaters. Though most anglers fish from a boat, two fishing piers offer great bank fishing opportunities for those without a boat.	
Bluegill	Excellent	There is a high number of 6-8 inch fish. The best fishing is in May, but bluegill can be caught year-round. Crickets and worms under corks during the spawn are great baits. Later in summer try the stake beds or deep (6'-10' water) brush piles. Catch bluegill in a feeding frenzy during the summer months where mayflies are hatching along the shoreline. Also try deeper rocky points during the summer with pieces of night crawlers.	
Channel catfish	Excellent	In the lake, the best fishing occurs during the late spring along chunky rock shorelines and rip-rap shorelines around the dam and marinas. Good fishing occurs along the river channel during summer and on flats along the secondary channels. Jug fishing in the embayments is often a good way to catch a mess of channel catfish.	
		In the tailwaters there is year-round fishing. However, blue catfish outnumber the channel catfish in the tailwater area. Two piers offer good bank fishing vantage points, while fishing from a boat might me best downstream around woody debris, stumps and creek mouths.	
Crappie	Good	After a few good spawns, anglers should notice a lot of crappie in the 10-13 inch range. Anglers fishing deep are catching fair numbers of keeper-size white crappie. Anglers fishing shallow along rocky shorelines are catching fair numbers of black crappie. Try casting toward the shoreline with a curly tail jig. Crappie are vulnerable during late winter or early spring as they move toward shallow water areas for spawning. Fish are shallow in early to mid April around buttonball bushes, brush piles and stake beds. Crappie move to secondary channels, drops and flats near creek channels during the fall and winter.	
Largemouth bass	Excellent	Several good year classes have resulted in some great fishing. There are a lot of bass in the 15 to 18 inch range. During the spring, fish move to wood structure along the shoreline. During the summer big fish are found along the main lake drops and secondary channels. Fishing the weed beds is good with top water baits on cloudy, rainy days during the summer. Smaller fish can be found in the backs of embayments around brush and chasing shad and minnows during late summer. During the fall, fishing around boat docks and rocky shorelines is good.	
		There is some fine bass fishing in the tailwaters below Kentuck Dam. Try fishing white spinner baits along the rip rap shoreline.	

Location/Species	Forecast	Comments		
KENTUCKY LAKE and TAILWATER (continued)				
Redear sunfish	Excellent	Best fishing is in late April and early May. Redear like areas with gravel substrate. If there are yellow mustard flowers inundated by water in the backs of embayments during this time, this is a good place to find redear. Submerged pondweed clumps also hold redear sunfish. These fish primarily feed on the bottom for bugs, snails and small mussels; hence you need to fish near or on the bottom. Redear generally spawn a little deeper than bluegill.		
Sauger	Fair	In the lake, moderate population of harvestable-size (≥14 inch) fish. Best fishing is in fall and winter along secondary channels. Try trolling crankbaits.		
		In the tailwaters there is a higher abundance of harvestable-size sauger. This is a good winter fishery. Boat fishing is ideal, though two new fishing piers offer good bank fishing opportunity.		
Smallmouth bass	Good	Best fishing along rocky shorelines and bluffs, creek channels at mouths of embayments, the canal main lake points, and main lake stumps. Night fishing during the summer months along gravel bars at the mouths of LBL embayments and rocky shorelines is good with dark color spinner bait or crayfish color creature baits.		
Spotted bass	Fair	Spotted bass are not normally targeted, but caught while fishing for largemouth or smallmouth bass. Spotted bass migrate to the streams at the backs of embayments during the early spring. Gravely bars and points are prime areas to fish during the summer. Secondary points are good during the fall.		
Striped bass	Fair	Striped bass and its hybrid are being caught from the lake while fishing for white bass. Trolling with 6-inch deep diver baits which imitate minnows is effective.		
		Better fishing is below Kentucky Dam in the tailwater. Fish congregate below the dam during periods of discharge through the generator section of dam. Live herring is the preferred bait. Most of the bank fishing has opened back up following bridge construction with the addition of two new fishing piers. From the shoreline, fish with 4-inch sassy shad type baits.		
White bass	Fair	Try fishing in the main lake during summer months when white bass are found along secondary creek channels near the main river channel. Also look for jumps when groups of white bass chase shad minnows near the water's surface. In-line spinners and silver spoons work great for catching white bass. Look for main channel points and bends; also near the canal.		
		Good numbers are also found in the Kentucky Dam tailwater area along with its cousin the yellow bass. When fishing from the bank or boat, try small curly tail jigs or in-line spinner type baits.		
Yellow bass	Excellent	Lots of fish but they don't get real big. Their taste is better than most anglers perceive. Casting the rocky shoreline with a curly tail jig or rooster tail works great. See a lot casting the rocky shoreline in the spring while crappie fishing.		

Location/Species	Forecast	Comments
KENTUCKY RIVER	255 miles (main	stem)
Bluegill*	Fair	Shoreline areas around woody debris out of the current hold some quality fish but in low numbers.
Channel catfish	Fair	Look for silt-free rocky habitat or below dams or palisades.
Common carp	Fair	Bottom fish with grain-based baits below dams.
Crappie*	Fair/Good	Currently good number of crappie in the Kentucky River; locate in backwater/creek-mouth areas with low current. Fish up to 12 inches or larger possible.
Flathead catfish	Fair	Large fish present around large boulders and log jams.
Hybrid striped bass	Fair	Sporadic catches of 15 to 20-inch fish below dams. Numbers are increasing due to a stocking program which began in 2006.
Largemouth bass*	Fair/Good	Good population in lower pools. Locate in cover just below dams, in creek mouths, bank slides, or around timber.
Muskellunge	Good	Best in spring around creek mouths, and in summer around timber and creek mouths. Stocked annually in upper river. More fish in upper half of river (pools 6-14).
Sauger	Good	Found around sand bars, creek mouths and below dams around rock and sand bars; numbers are increasing due to a stocking program which began in 2006.
Smallmouth bass	Fair	Nice fish possible in mouths of quality tributaries. Also found around rock piles and walls.
Spotted bass	Fair	Sporadic catches of small to medium-size fish; usually below dams but also around palisade rock wall areas.
Walleye	Fair	Present in low numbers due to past stockings and escapement from reservoirs in the headwaters; good-size fish possible below dams.
White bass	Fair/Good	Many 10-12 inch fish below dams; numbers are increasing due to a stocking program which began in 2006.

^{*}These are considered "standing water species" and do best in the river after prolonged periods of low-flow conditions. High water or flooding conditions for extended periods will reduce their numbers in the Kentucky River.

KINCAID LAKE	183 acres	(Pendleton County)
Bluegill	Poor	Abundant, most fish are less than 6 inches.
Channel catfish	Good	Catchable-size stocked each year; good potential for fish over 15 inches.
Crappie	Fair	Abundant; most between 7-9 inches.
Largemouth bas	ss Exceller	Very good numbers of fish over the 12-inch size limit; good numbers of fish over 15 inches; excellent potential for a trophy fish. Note: 12-inch size limit.

_ocation/Species	Forecas	t Comments
LAKE CARNICO	114 acres	(Nicholas County)
Bluegill	Poor	Numbers are low, however a few fish greater than 6 inches can be found.
Channel catfish	Fair	Fair numbers of flathead and channel catfish can be found. Lake is stocked with channel catfish.
Largemouth bass	Fair	Good population of bass; good numbers of fish greater than 15 inches and a few 20-inch bass are present. Fifteen hundred 4.5-inch largemouth bass were stocked in the fall of 2011 and 900 during 2012 to help offset effects of poor spawns in those years. Aquatic vegetation is present at the lake so bass anglers my want to concentrate their efforts near weed beds.
Redear sunfish	Poor	Numbers are low, however a few fish greater than 5 inches can be found.
LAKE MALONE 7	67 acres	(Muhlenburg, Todd, and Logan counties)
LAKE MALONE 7 Bluegill	67 acres Fair	(Muhlenburg, Todd, and Logan counties) Population mostly comprised of 4-6 inch fish. Some larger fish available.
Bluegill	Fair	Population mostly comprised of 4-6 inch fish. Some larger fish available. Stocked annually, good number of 14-16 inch fish with a few 20-inch plus available.
Bluegill Channel catfish	Fair Good	Population mostly comprised of 4-6 inch fish. Some larger fish available. Stocked annually, good number of 14-16 inch fish with a few 20-inch plus available. Moderate number of 9-10 inch fish. Fish brush piles and fallen trees along some of the steeper banks.

ocation/Species	Forecast	Comments
AKE REBA 76 ac	res (Madisor	n County)
	gram for chemic	be a problem within the lake, making bank fishing difficult. KDFWR personnel have all and biological removal of the vegetation. In 2007, 2010 and 2013 grass carp were se fish is illegal.
Bluegill	Good	Across the board all numbers went up in 2013; most relevant are the numbers of fish over 6 inches. This lake is making gradual improvements in terms of the bluegill fish as we are making gradual improvements in the aquatic vegetation. Anglers should expect a slightly better year this year. Great lake to introduce a kid to fishing with the excellent bank access.
Channel catfish	Good	Stocked annually. Sampling shows high numbers of these stock-size fish and sever over this size. Note: 12-inch size limit; limb lines (set lines), jug lines and trot lines are illegal to use at this lake.
Largemouth bass	Good	Number of fish sampled over the 15-inch minimum size limit went up significantly las year but there are still few fish over 20 inches. Largemouth bass fishing should rema good. Reports from anglers are that this is a better summer lake than a spring lake there is a need to get away from the crowds along the banks to find the larger fish. Note: 15-inch minimum size limit.
Redear sunfish	Fair	Fair numbers within the lake, most less than 7 inches long; see bluegill.
AKE WILGREEN	169 acres (N	Madison County)
Blue catfish	Good	Stocked annually since 2009. Fishing success should be improving in the next few years. Can be difficult to determine from channel catfish with the main difference be the anal fin (straight in the blues and rounded in the channels).
Bluegill	Good	Similar number as previous years and most fish are less than 6 inches. The popular is healthy up to that point. Should give good fishing opportunities for 2013, but noth like the Wilgreen of several decades ago.
Largemouth bass	Excellent	High numbers of fish over 15 and 20 inches. Larger fish are fat and healthy for their size. All indicators are continuing to show increase in this population. Should be another great year.
Redear sunfish	Good	Similar numbers to previous years and most fish are less than 7 inches; see bluegill
AUREL CREEK RES	SERVOIR 43	acres (McCreary County)
Channel catfish	Fair	Stocked every other year.
Largemouth bass	Good	Numerous 12-14 inch fish with a few fish larger than 15 inches.

Location/Species	Forecast	Comments
LAUREL RIVER LAKE	5,830 acres	(Whitley and Laurel counties)
Bluegill	Good	Fairly abundant 6-7 inch fish. Try crickets along rock walls in upper Laurel River Arm during summer months.
Crappie	Fair	Still not great for numbers, but some large crappie are available to skilled anglers. Upper Laurel River near city dam is best area.
Largemouth bass	Good	The largemouth population has steadily gotten better over the last decade with improved numbers of 14-17 inch fish and some over 20 inches. Upper Laurel River arm is best area.
Rainbow trout	Fair	Stocked annually in January. Should be adequate numbers of 10-14 inch trout and a few larger fish. Try trolling small cranks or spoons during winter and early spring for good catches of stocked fish.
Smallmouth bass	Good	Good winter fishery with numbers of fish over 3 pounds and a few over 6 pounds. Smallmouth fishing at Laurel can be difficult at times but persistent anglers have a chance of catching a trophy fish. Try casting jigs or swim baits in winter and early spring for best success.
Spotted bass	Good	Stable population of mostly 11-14 inch fish with a few exceeding 15 inches. Best in the mid and lower sections of the lake. Look for schooling fish chasing shad on the surface in early fall.
Walleye	Good	Several year classes contributing to fishery. Most of the larger fish (20-26 inch) from strong 2009 year class. Try surface baits at night during spring, or troll deep diving crank baits or spinner rigs across points and islands in summer and fall.
White bass	Poor	A consistent fishery has failed to develop, although a few can still be caught each year. Best numbers in Laurel River arm.
LIBERTY LAKE 81	acres (Casey	County)
Largemouth bass	Fair	Moderate numbers of mostly smaller fish (<13 inches), although fish larger than 18 inches are available.
Spotted bass	Poor	Low density of mostly smaller fish (<12 inches).
MARION COUNTY LA	(E 32 acres	(Marion County)
		ember) oxygen levels are too low to support fish below 11 feet.
Bluegill	Good	Good numbers of 6-inch plus fish.
Channel catfish	Good	All sizes available. Higher than average numbers of 12-inch plus fish available.
Largemouth bass	Fair	High numbers of 11-13 inch fish.
Redear sunfish	Fair	8-inch plus fish available. Larger fish up shallow in early-mid April for spawn then move out to thermocline depth.
White crappie	Fair/Good	Large fish available; fish laydown trees in the upper end of the lake in spring and brushpiles in late winter and summer.

Location/Species	Forecast	Comments	
MARTINS FORK LAKE 334 acres		(Harlan County)	
Bluegill	Fair	Good numbers. Most harvested fish are 7.25 inches or less. Fish shallow mud flats in spring or summer.	
Channel catfish	Good	Most fish are less than 20 inches; fingerlings were stocked in 2010 and 2012. Fish at night for better results as water is very clear. There are several good bank fishing areas. Martins Fork Creek entering the lake when murky can have good numbers of fish.	
Crappie	Fair	White crappie are present. Harvested fish are typically 9 to 10 inches long. Occasional large fish up to 12-13 inches. Crappie are scattered in shallow water in spring, and congregate in tight groups in standing timber in summer.	
Largemouth bass	Fair	Excellent growth but low recruitment and low numbers. A good year class of fish spawned in 2010 should improve numbers in coming years and fingerlings were stocked in 2011. Numbers over 15 inches are above average at present for the lake. This lake contains 4 species of black bass that may be caught by anglers (largemouth, smallmouth, spotted and coosa bass). Largemouth bass numbers account for approximately 80% of the black bass in the lake.	
Walleye	Poor	The last stocking of Erie strain walleye occurred in 2004. Very few of these fish remain This fishery is currently rebuilding with the first stocking of native strain walleye in 2013 Keeper-size fish will not be common until 2015 fishing season. There will be a few keeper-size fish available late in 2014.	

MAUZY LAKE 81	acres (Union Co	ounty, Higginson-Henry WMA)
Bluegill	Fair/Good	Large number of 3-5 inch bluegill, but number of 6-8 inch fish in the population is increasing.
Channel catfish	Good	Stocked annually; many 15-22 inch fish in the population.
Largemouth bass	Good/Excellent	Number of 12-14 inch and 15-20 inch fish have declined over the last couple of years. An increased number of smaller fish in the population should bring these numbers back up as they grow into the upper size ranges in the next few years. Fish plastic baits and spinner baits around hardwood brush piles, especially in spring/early summer.
Redear sunfish	Fair/Good	Redear sunfish population continues to improve annually. Number of fish over 8 inches continues to increase.

MCNEELY LAKE	51 acres (Jeffers	son County)
Bluegill	Good	Good numbers of fish through 8 inches.
Channel catfish	Fair/Good	Catchable-size stocked each year; large fish possible.
Common carp	Fair	A few large fish present.
Crappie	Fair	Quality size fish possible; fish near weed beds and fallen trees, particularly in spring.
Largemouth bass	Fair/Good	Good numbers of fish over the 12-inch size limit; best cover is water willow, fallen trees and stumps.
Redear sunfish	Excellent	Good numbers of fish through 9 inches; ≥10-inch fish are present.
Warmouth	Fair	Large fish present around weed beds.

Location/Species	Forecast	Comments	
METCALFE COUNTY I	LAKE 22 ac	cres (Metcalfe County)	
		ember) oxygen levels are too low to support fish below 7 feet.	
Bluegill	Fair	High numbers overall with fair numbers of 6-inch fish; however, few exceed 7 inches.	
Blue catfish	Fair	Fair numbers overall, but some trophy-size fish (30-inch plus) available. Fish with live or cut bait near creek channel. Stocked annually since 2009.	
Channel catfish	Fair/Good	Fair numbers available. Stocked every other year.	
Crappie	Poor/Fair	Smaller sized (8-9 inch) and not overly abundant.	
Largemouth bass	Good/Excellent	Fair numbers of 15-inch plus fish. Trophy fish (20-inch plus) potential consistently above average. Use trout colored jerkbaits in May.	
Rainbow trout	Fair	600 fish stocked in March for a put and take fishery. Water too warm by mid-May to support trout.	
MILL CREEK LAKE	109 acres (M	lonroe County)	
Summertime (mid-Ju	ine through Septe	ember) oxygen levels are too low to support fish below 12 feet.	
•		, ,,	
Blue catfish	Fair	Stocked annually since 2008 with 550 fish per year.	
Bluegill	Good	Good numbers of 6-inch plus fish.	
Channel catfish	Fair	Stocked every other year.	
Crappie	Fair	Fair number of 8-9 inch fish. Larger crappie present but at low numbers.	
Largemouth bass	Good/Excellent	Good numbers of larger fish (15-inch plus) overall; 18-inch plus fish numbers well above average.	
Redear sunfish	Fair	Lower numbers overall, but large fish (10-inch plus) available.	
MILL CREEK LAKE	41 acres (Po	well/Wolfe County – Natural Bridge State Park)	
Bluegill	Good	Better numbers recently; high numbers of fish over 6 and 8 inches. Water is extremely clear and can be difficult to fish.	
Channel catfish	Good	Stocked annually. Note: limb lines (set lines), jug lines and trot lines are illegal to use at this lake.	
Largemouth bass	Good	Similar numbers to previous years. Holds nice 20-inch plus fish but this lake can be difficult to fish. The timber areas in the upper reaches are the best bet. Note: 15-inch minimum size limit.	
Rainbow trout	Excellent	Stocked annually. Best time to fish is within the first few days after stocking.	

Old and New Kingfisher lakes are in the process of being drained and renovated. The lakes will be allowed to dry throughout the summer of 2014 and the renovation will occur during the fall of 2014. To see when the lakes will be available for fishing check the KDFWR website at http://fw.ky.gov

(Daviess County)

21 acres

NEW KINGFISHER LAKE

Location/Species	Forecast	Comments
NOLIN RIVER LAKE	5,795 acres	(Grayson, Edmonson and Hart counties)
Catfish	Good	<u>Channel catfish</u> - Good number of 15-20 inch fish and many larger. Fish upper lake (Sportsman's Paradise and Cane Run) and river area in early spring and rocky main lake banks mid-May through June. <u>Flathead catfish</u> - lower numbers than channels but several large (30+ lb.) fish harvested annually. Fish late May through June along the steeper rocky banks.
Crappie	Good	Crappie angling should continue to be good at Nolin in 2014 with a good number of 9-11 inch fish available for harvest. During summer months fish deeper brush piles 18-20 feet deep mid-lake for larger fish.
Largemouth bass	Good	Many 15-20 inch largemouth in the population. Anglers should benefit from a good number of harvestable bass in the population in 2014.
Walleye	Fair	Most fish 13-16 inches with a few up to 20-21 inches. During early spring fish Bacon Creek and above with lead head jigs and plastic grubs. In the summer, troll Sportsman's Paradise, Conoloway Creek and the longer main lake points in the Red Rock and Rock Creek areas.
White bass	Excellent	Large number of 13-15-inch white bass in the population. Fish Bacon Creek and upper river during spring spawning run with lead head jigs and grubs. Fish longer main lake points or jumps in summer. Also in summer fish at night under lights near main channel with jigging spoons or live minnows.
OHIO RIVER		
Blue catfish	Good	Good populations in tailwaters and deep holes around outside bends.
Channel catfish	Excellent	Look for silt-free, rocky habitat and embayments.
Common carp	Fair	Large fish possible. Tributaries and backwaters away from current are best areas. Increasing numbers of silver carp and bighead carp.
Crappie	Good/Excellent	Backwater areas and creek mouths with brushy habitat are best producers.
Flathead catfish	Good	Potential for very large fish; look for cover such as sunken logs, boulders or other large objects.
Hybrid striped bass	Excellent	Fish up to 6 pounds in and around tailwaters.
Largemouth bass	Good	Tributaries and backwaters away from current are best areas.
Sauger	Good	Spring and fall below dams with minnows or jigs on bottom.
Smallmouth bass	Fair	Low population in mid/upper river in sections of rocky habitat and tailwaters.
Spotted bass	Fair	Fish the main channel around cover.
Striped bass	Fair	Mid-summer to late fall good for 6-10 pound fish.
Walleye	Fair	Occasionally found in tailwaters.
White bass	Good	Good population.

Location/Species	Forecast	Comments
PAINTSVILLE LAKE	and TAILWATER	1,139 acres (Morgan and Johnson counties)
		Boaters are asked to take measures to prevent the spread of this damaging non-native g and disinfecting boats, trailers and equipment.
Channel catfish	Fair	Large catfish present in low densities in upper lake. Down-lake, large fish are often located deep; fish that are shallow can be found around rock riprap shoreline areas.
Crappie	Good	Most of successful fishing is done on the upper end of the lake into Open Fork and Little Paint Creeks; fish sampled up to 15 inches. Late March can often provide good catches of keeper fish in shallow water near Patoker boat ramp. Look for sunny days at this time of year for best fishing. Lower lake will have larger fish often holding at edges of hydrilla beds.
Largemouth bass	Fair	High survival of young fish due in part to presence of hydrilla which offers ample cover. Stable population with high numbers of fish <12 inches; fish sampled up to 24 inches; slow growth rates; number of fish over 15 inches is increasing. March is popular for catches of lunker fish.
Rainbow trout	Good	Good fishing present in shallow water during winter/early spring; fair fishing present in summer at depths of 20-25 feet. The catch and release portion of the tailwater can offer good quality rainbow and brown trout fishing.
Smallmouth bass	Poor	Poor spawns; fish sampled up to 22 inches (lower lake areas seem best for larger fish) 18-inch minimum size limit implemented in 2005; changes made in water releases at dam in 2006 to improve summer cool water habitat; spawning benches being placed in lake to aid recruitment. Supplemental stockings of native strain fish have occurred from 2009-2012.
Spotted bass	Poor	Numbers are down; some large spotted bass present up to 14 inches; best fishing in lower half of lake.
Walleye	Fair	Fair numbers through 27 inches; potential for trophy size; look for active fish to be suspended in March; also May and October are good months to fish with live night crawlers on a light jighead fished off the bottom. During summer target weedbed edges and dropping minnows or nightcrawlers down these edges. Last several year classes are doing well and will provide quite a few keeper fish for 2014.
White bass	Poor	Low numbers but good size fish. Fish will congregate in Little Paint and Open Fork arms in March.

Location/Species Forecast Comments
PAN BOWL LAKE 98 acres (Breathitt County)

Eurasian water milfoil is present in Panbowl Lake. Boaters are asked to take measures to prevent the spread of this damaging non-native invasive species by properly inspecting and disinfecting boats, trailers and equipment. Grass carp were stocked in Panbowl Lake in 2010 to control the Eurasian water milfoil. All grass carp caught in Panbowl Lake must be released immediately.

Bluegill	Excellent	Excellent numbers and size distribution up to 7.5 inches throughout the lake. Fish early or late in the year when milfoil is not a problem.
Channel catfish	Good	Most fish are less than 20 inches; fish were stocked in 2010 and 2012; lots of shoreline areas for bank fishing.
Crappie	Fair	Black and white crappie are present; fish sampled up to 12 inches; most are less than 9 inches; tend to congregate in tight groups in deeper areas of the lake.
Largemouth bass	Poor	Numerous fish up to 12 inches and good numbers of fish above 18 inches; lake contains a lot of aquatic vegetation providing cover for large fish. These fish see a lot of angling pressure often making success for large fish tough.
Redear sunfish	Excellent	Excellent numbers and size distribution up to 8.5 inches. May and June are good months to fish around weed beds and near docks.

PENNYRILE LAKE	47 acres	(Christian County)
Bluegill	Fair	Some nice bluegill are present as a result of the stunted bass population. Numerous fish less than 6 inches also available.
Channel catfish	Good	Good numbers of 12-15 inch catfish and a few larger individuals present as well. Fishing around fallen trees and beaver huts with night crawler or cheese bait should provide success.
Crappie	Poor	Low overall numbers of crappie, but they are large fish.
Largemouth bass	Fair	A quiet and scenic lake with high numbers of bass 12 inches and smaller. It is a great place to take kids fishing! Bass are protected between 12-15 inches. Harvesting some smaller ones is allowed, and will help the population.
Redear sunfish	Fair	Numerous fish in the 7-9 inch range. Few redear present greater than 10 inches. Try fishing a small piece of night crawler under a cork near the bottom for best success.
REBEL TRACE LAK	E 19 acres	(Menifee county)

Watershield has now replaced the Eurasian watermilfoil and is covering about 80% of this lake; therefore fishing may be difficult, especially in the summer and fall. This lake is also rapidly filling in. Due to federal ownership of this lake the use of aquatic herbicides is prohibited at this time.

Bluegill	Good	Fair numbers of 6-inch bluegill are present with occasional 8-inch fish caught. Fish shallow shoreline areas and downed timber in spring. Also, as the lake fills in, numerous spawning beds can be found off shore in shallow areas where the aquatic vegetation is less dense.
Largemouth bass	Poor	Lake receives heavy pressure therefore fewer numbers of bass are caught. Though limited, fish up to 19 inches can be found. Fish woody structure and weed beds.
Redear sunfish	Good	Good population of 7-inch fish with some 9-inch fish present.

Location/Species	Forecast	Comments						
ROUGH RIVER LAKE and Tailwater		5,100 acres (Breckinridge and Grayson counties)						
Catfish	Good	Good number of channel catfish in the 20-25 inch range. Lower number of flathead catfish, but trophy fish available. Tailwater good in early summer and again during fall drawdown. Fish upper lake areas in early spring and steeper rocky banks during mid May-June for spawning fish.						
Crappie	Good	Large increase in number of 7-8 inch fish due to very successful spawn in 2011. Still good number of 9-11 inch fish in the population with 12-14 inch fish available. Fish deeper (16-18 ft.) brush piles in mid-lake during summer for larger fish. Tailwater area is a good early spring crappie fishery.						
Hybrid striped bass	Excellent	Excellent population of hybrid striped bass. Many 20-inch plus fish available. Fish upper lake areas in spring. Troll main lake points in summer or fish jumps in main lake (Dam to Tules Creek on north fork and Dam to Little Clifty Creek on south fork). Fish main lake points and sand/mud flats in fall and winter. Tailwater area good spring fishery (April).						
Largemouth bass	Good	The number of 15-20 inch bass continues to be higher than it has for the last several years. Bass anglers at Rough should continue to enjoy a good number of harvestable size bass for the next few years.						

SHANTY HOLLOW LAKE 135 acres (Warren and Edmonson counties)

Summertime (mid-June through September) oxygen levels are too low to support fish below 10 feet.

Bluegill Fair Numbers of 6-inch plus fish up from last couple of years. Fair/Good Channel catfish All sizes available. Stocked annually. Fair Good numbers of 12-14 inch fish. Largemouth bass Redear sunfish Fair Fair numbers of 8-inch plus fish. Larger fish up shallow early in the spring then out to thermocline depths by early summer. White crappie Fair/Good Large fish (10-14 inches) available. Fish stakebeds and shallow brushpiles in spring; deeper brush in late winter and summer.

SMOKY VALLEY LAKE 36 acres (Carter County – Carter Caves State Park)

Fair

Bluegill

Due to slow growth of the largemouth bass population, a new regulation was established for the 2014 fishing season. Largemouth bass will now be under a 12-15 inch slot limit. All fish between 12 and 15 inches must be immediately released. There will be a 6 fish daily creel limit with only 1 fish in the daily creel allowed over 15 inches. KDFWR anticipates improvements in the largemouth bass population under this new regulation.

Slightly better than last season; most fish less than 7 inches

Diacyiii	ı alı	olightly better than last season, most list less than 7 mones.
Channel catfish	Good	Stocked annually. Note: limb lines (set lines), jug lines and trot lines are illegal to use at this lake.
Largemouth bass	Fair	There are plenty of chances to catch a good number of fish, but the size is still not there for this population. Note: 12-15 inch slot limit; 6 fish daily creel limit (only 1 fish in the daily creel allowed over 15 inches).

Location/Species	Forecast	Comments
SPURLINGTON LAKE	36 acres	(Taylor County)
Summertime (mid-Ju	une through Sept	rember) oxygen levels are too low to support fish below 7 feet.
Bluegill	Fair	Fair numbers of 6-inch plus fish.
Channel catfish	Good	All sizes present. Stocked annually.
Largemouth bass	Excellent	Better than average numbers of larger fish (15-inch plus). Big bass (20-inch plus) opportunities above average.
Redear sunfish	Fair	Fair numbers of larger fish (8-inch plus) available. Occasional 10-inch fish available.
STANFORD RESERVO		(Lincoln County)
Largemouth bass	Good	Good density of 12-16 inch fish with some fish larger than 18 inches available.
TAYLORSVILLE LAKE	3,050 acres	(Spencer, Anderson and Nelson counties)
Blue catfish	Good	Population stable; stocked annually since 2002. Fish with live bait, cut bait or crank
		baits (open water areas). Increasing number of fish over 30 inches.
Bluegill	Good	Good number of fish 6 inches and longer around standing timber in coves.
Channel catfish	Good	Many fish of all sizes. Look for bigger fish in the timber in spring, heads of creeks in fa and points next to channel any time.
Crappie	Fair/Good	Good number of fish at and above the 9-inch size limit. Great white crappie spawn in 2013. Note: daily creel limit 15 fish.
Hybrid striped bass	Fair	Overall numbers lower, but fair numbers of fish greater than 15 inches. In spring and fall, there is good fishing in headwaters; look for jumps early and late in day July-September.
Largemouth bass	Good	Very good numbers of 12 to 15-inch fish; Good number of 15 to 18-inch fish; best in timber coves; try embayments in spring and fall. Fish over 20 inches possible.
White bass	Poor	Low numbers in Salt River above the lake in spring and fall; fish mostly range from 10-13 inches.

Location/Species	Forecast	Comments
THREE SPRINGS LA	KE 30 acres	(Warren County)
Lake is very shallow	(no deeper tha	n 6-7 ft)
Bluegill	Fair	Fair numbers of 6-8 inch fish.
Channel catfish	Good	Good numbers and good size fish (15-inch plus) available. Stocked periodically through the spring and summer.
Largemouth bass	Fair	Numerous fish available; few fish larger than 12 inches. Fish trout colored swim baits in May.
Rainbow Trout	Good	Stocked periodically through winter and early spring months (November – March) for put and take fishing. Lake too warm for trout by early-mid May.
Redear	Fair	Fair number of 9-10 inch fish available.
WASHBURN LAKE	18 acres (O	hio County)
Bluegill	Fair	Low number, but many fish in 7-inch range.
Channel catfish	Good	Stocked annually. Good number of fish in the population, many 15-20 inches.
Largemouth bass	Poor	Population dominated by slower growing sub-legal fish. Few fish greater than 12 inches.
Redear sunfish	Fair	Low number, most fish in the 5-6 inch range, but some 7-8-inch fish available.
WEST FORK DRAKES	S RESERVOIR	88 acres (Simpson County)
Summertime (mid-J	une through Sep	otember) oxygen levels are too low to support fish below 8 feet.
Blue catfish	Fair	350 fish stocked annually since 2006.
Bluegill	Fair	Fair numbers of 6-inch plus fish. Few 8-inch plus fish.
Channel catfish	Good	Numerous sizes available. Fishing best at night near creek channel.
Largemouth bass	Fair/good	Better than average numbers of 12-inch plus fish due to good spawn in 2010.
Muskie	Fair	Fishing best in the lower half of the lake near vegetation beds in winter to early spring. Fishing good below the dam year-round using large inline spinners and jerk baits.
Redear sunfish	Good	Better than average number of 6-inch plus fish.
WOOD CREEK LAKE	672 acres	(Laurel County)
Bluegill	Fair	Adequate numbers of 6-7 inch bluegill.
Channel catfish	Good	Stocked every other year. Consistently produces 12-18 inch fish and remains one of the better catfish fisheries in this area of the state.
Crappie	Fair	Good density of smaller fish with relatively few fish exceeding 9 inches.
Largemouth bass	Good	Numbers are slightly down in recent years but the population still has good size range with some fish exceeding 20 inches.
Rainbow trout	Fair	Stocked annually in October and February.

Location/Species	Forecast	Comments
WOOD CREEK LAKE	(continued)	
Redear sunfish	Fair	Moderate density of 8-inch and larger redear are available. Mid and lower lake areas are best.
Spotted bass	Fair	Most are less than 12 inches. Mid and lower lake areas are best.
YATESVILLE LAKE	2,314 acres	(Lawrence County)
Bluegill	Excellent	Excellent numbers and size distribution throughout the lake; numerous fish 7.0-7.5 inches.
Channel catfish	Excellent	Quality size distribution and densities of catfish exist in all areas of the lake; recent stockings have greatly increased numbers; during fall to early winter many fish are shallow in upper lake; popular lake for jugging and limb lines.
Crappie	Good	Good numbers of 8-15 inch fish. Traditional brushy areas and deadfall trees are good for spring spawning fish, with points from shallow mudflats to deeper water being good in the fall. This fishery consists only of white crappie.
Flathead catfish	Fair	Good size distribution with fair numbers; rocky shoreline and brush will have better numbers. Recent years have had shown good survival of young fish.
Largemouth bass	Good	Good size distribution of fish with good numbers of 15-22 inch fish. Supplemental stockings of fingerling bass were done in 2009, 2012, and 2013 to aid average recruitment. Popular lake for tournament fishing as most of the year produces good results. Check out KDFWR web page for GPS coordinates of submerged fish habitat structures.
Redear sunfish	Fair	Population is building in numbers; annual stockings of fish from 2010-2013; larger fish in 2012 were up to 8 inches.

Fishing Tips for Kentucky and Barkley Lakes

Largemouth bass

In the spring, try a Rapala #11 or #13 gold with black back or floating or shallow diving stick baits. Also try black/brown or black/blue ½ ounce jigs with #11 or #1 pork frog in the yellow mustard flowers in the backs of embayments. Around woody structure use a spinner bait or plastic lizards in either pumpkin or june bug color.

During summer, fish ledges near river channels with Carolina-rigged and Texas-rigged soft plastics and deep-diving crank baits. Also fish stumps on the main lake in the mornings and evenings with topwater baits like the zara spook. Current is the key to these main lake stumps being productive. Using a 10-inch plastic worm fished along the ledges during the summer months is also good. A good color is plum.

During fall, fish along main lake ledges around stumps and brush with shad or blue/chartreuse-color crank baits. Use top water baits (zara spook or devil's horse) in the flats of embayments. Use white spinner baits around main lake stumps.

The recent dry summers have really brought on the aquatic vegetation. The beds of weeds are extremely productive during the late spring, summer and early fall. Top water baits such as spooks, stick baits, plastic frogs, and buzz baits are great choices. Plastics, such as worms, and spinner baits are also good when fishing the edges of the weed beds.

Crappie

In spring, fish shallow and over flats with jigs and minnows. Cork fish around buttonball bushes. During summer, white bass anglers trolling small crankbaits near the river channel and primary points in the lake catch many crappie. During fall, fish jigs and minnows next to deep ledges formed by the river channel. Fish may also be caught suspended on flats adjacent to the secondary channel around 10 feet deep near structure. Fall is also a good time to catch black crappie along gravelly shorelines fishing with a curly tail jig.

Blue catfish

During the summer, fish along the main river channel with minnows, cut shad/skipjack herring and stink baits.

Channel catfish

In spring, fish with leeches or night crawlers around rip rap. During summer, fish with cut shad, chicken livers, minnows, leeches and worms along the main river channel.

Sauger

Troll main-lake bars during the late summer and early fall with wiggle-warts or crankbaits which bump the bottom. During fall and winter, use jigs and minnows along the old river channel, along primary points near the river channel, and where a secondary channel meets the old river channel. A good site for fishing is along the river channel just north of the Paris Landing Bridge (US Highway 79).

White bass

Fish the jumps from July-September with slab spoons or white roostertails.

Fishing Tips for Kentucky and Barkley Tailwaters

Blue and channel catfish The best time is in May-June using cut bait or live bait just off the bottom.

Sauger Fish ¾-oz or 1-oz jigs with minnows bumping bottom in back eddies and

near current. Trolling is also good during the winter and early spring.

Best fishing is between December and February.

Striped bass and hybrids The best time is from April-June using live bait (skipjack herring) or 4-

inch sassy shad baits with a 1-2 ounce head in pearl or chartreuse color. The amount of weight depends on the current and should allow baits to bump the bottom. Live bait can be caught in the tailwaters using a cast

net or a skipjack rod.

Fishing Forecast Cheat Sheet (Summary of lakes forecasting good or excellent fishing for each species listed)

	Largemouth	Smallmouth	Spotted		Redear		Rainbow	Brown	
Water body	bass	bass	bass	Bluegill	sunfish	Crappie	trout	trout	Muskellunge
Barkley Lake/tailwater	_			_		_			
Barren River									
Barren River Lake	_			_		_			
Beaver Lake									
Beshear Lake									
Boltz Lake				_					
Briggs Lake									
Buckhorn Lake/tailwater									_
Bullock Pen Lake	_			·					
Carpenter Lake	·								
Carr Creek Lake			1/						
Cave Run Lake			<u> </u>						
Cedar Creek Lake			•						<u> </u>
Clear Creek Lake	•			-		•			
Corinth Lake				•					
Cumberland Lake		-/	./						
Cumberland Tailwater	*	₩				V	_/	-/	1
Dale Hollow Lake			./			†	V		†
Dewey Lake				./		./			†
Elkhorn Creek				V		V			+
Elmer Davis Lake		V	 		./	 			+
Fagan Branch Lake	V		 			 			+
Fishpond Lake			 	./	V	 	./		+
Fishtrap Lake/tailwater	V		 	V		. /	V		+
Floyds Fork Creek				V		V			+
-							V		+
Grayson Lake			-	V					
Green River		V							
Green River Lake						V			
Greenbo Lake							✓		
Guist Creek Lake									
Herrington Lake									
Kentucky Lake/tailwater		√		√	√	✓			
Kentucky River									✓
Kincaid Lake									
Lake Malone									
Lake Reba									
Lake Wilgreen	✓			\checkmark	\checkmark				
Laurel Creek Reservoir									
Laurel River Lake	\checkmark	\checkmark	\checkmark						
Marion County Lake				\checkmark					
Martin's Fork Lake									
Mauzy Lake	\checkmark								
McNeely Lake				\checkmark	\checkmark				
Metcalfe County Lake									1
Mill Creek Lake (Monroe Co.)				✓					
Mill Creek Lake (Powell/Wolfe Co.)	\checkmark			\checkmark			\checkmark		
Nolin River Lake						√ .			
Ohio River	\checkmark					√			
Paintsville Lake/tailwater						V	\checkmark		
Panbowl Lake				$\overline{}$	V				
Pennyrile Lake									
Rebel Trace Lake									
Rough River Lake					•	_			
Smoky Valley Lake									
Spurlington Lake									
Stanford Reservoir	_								
Taylorsville Lake	—			_					
Three Springs Lake									
Washburn Lake							*		1
West Fork Drakes Reservoir					1/				1
West Fork Diakes Reservoir			•		¥	•			
Wood Creek Lake									

^{**} Additional species can be found on following page.

Fishing Forecast Cheat Sheet (Summary of lakes forecasting good or excellent fishing for each species listed)

	Striped	Underid atripod	White			Channel	Divo	Flathead
Water body	bass	Hybrid striped bass	bass	Walleye	Sauger	catfish	Blue catfish	catfish
Barkley Lake/tailwater		1	5000	Transys	Oddgoi	l ./	- Cathon	James
Barren River	 					V		
Barren River Lake	1	./				-/	./	./
Beaver Lake	 	V				<u> </u>	V	V
Beshear Lake	1							
Boltz Lake	 					V		
Briggs Lake	1					_		1
Buckhorn Lake/tailwater	+					<u> </u>		./
Bullock Pen Lake	1					V		V
Carpenter Lake	1							
Carr Creek Lake	1							1
Cave Run Lake	1			V				
Cedar Creek Lake	1	+	V	1		V		1
Clear Creek Lake	1					1		
Corinth Lake	1							
	 							
Cumberland Lake						V		
Cumberland Tailwater	1 	+						
Dale Hollow Lake	 	+		1				
Dewey Lake	┨ ├───	1		1		V	V	V
Elkhorn Creek	┨ ├───	1		1		V		
Elmer Davis Lake	 			1				
Fagan Branch Lake	 	1		ļ		√		
Fishpond Lake				1		,		
Fishtrap Lake/tailwater		\checkmark				\checkmark		\checkmark
Floyds Fork Creek								
Grayson Lake								
Green River		\checkmark				<u> </u>		\checkmark
Green River Lake						<u> </u>		
Greenbo Lake								
Guist Creek Lake						\checkmark		
Herrington Lake								\checkmark
Kentucky Lake/tailwater						\checkmark	$\sqrt{}$	
Kentucky River					\checkmark			
Kincaid Lake								
Lake Malone								
Lake Reba								
Lake Wilgreen							_	
Laurel Creek Reservoir								
Laurel River Lake				_				
Marion County Lake								
Martin's Fork Lake								
Mauzy Lake						V		
McNeely Lake								
Metcalfe County Lake								
Mill Creek Lake (Monroe Co.)						İ		
Mill Creek Lake (Powell/Wolfe Co.)								
Nolin River Lake								_
Ohio River					_/	7		
Paintsville Lake/tailwater		*			· ·	<u> </u>		•
Panbowl Lake	1					./		
Pennyrile Lake	1					-		
Rebel Trace Lake	1					V		
Rough River Lake	1	./				_		1
Smoky Valley Lake	1	V						
Spurlington Lake	1	+		1				
Stanford Reservoir	 	+		1		V		
	 	+		1				
Taylorsville Lake	 	+		1		V	V	
Three Springs Lake	┨ ├───	1		1		V		
Washburn Lake	↓	1		1				
West Fork Drakes Reservoir	 	1				V,		
Wood Creek Lake	 	1						
Yatesville Lake				İ]	✓		l